





Entrez	PubMed	Nucleotide	Protein	Genome	Structure	MIMO	PMC	Journals	Book
Search	PubMed	for					Go	Clear	
		Limits	Previe	w/index	History	Clip	board	Deta	ils
About Entr	'ez	Display Abs	tract		Show: 20	Sort		Send to Te	

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy

E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher **Batch Citation Matcher** Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

h

1: Nature. 2000 Sep 14;407(6801):242-8.

Related Articles, L

nature

Vascular-specific growth factors and blood vessel formation.

Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J.

Regeneron Pharmacueticals, Inc., Tarrytown, New York 10591, USA.

A recent explosion in newly discovered vascular growth factors has coincided with exploitation of powerful new genetic approaches for studying vascular development. An emerging rule is that all of these factors must be used in perf harmony to form functional vessels. These new findings also demand reevaluation of therapeutic efforts aimed at regulating blood vessel growth in ischaemia, cancer and other pathological settings.

Publication Types:

- Review
- Review, Tutorial

PMID: 11001067 [PubMed - indexed for MEDLINE]

Display Abstract	Show:	20	ž	Sort 👻	Send to Text
------------------	-------	----	---	--------	--------------

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09

C

cb h g fcg c b b







Go

Entrez

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Clear

Books

Search PubMed

Limits

for AL-2 AND Eph Preview/Index

History

Clipboard

Details

No items found.

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial

New/Noteworthy E-Utilities

PubMed Services Journals Database

MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries

LinkOut Cubby

Related Resources

Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

> Write to the Help Desk NCBI | NLM | NIH
> Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

> > Oct 13 2004 06:44:09







Go

Entrez

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed

• for AL-2 AND ephrin Limits

Preview/Index

History

Clipboard

Clear

Details

No items found.

About Entrez

Text Version

E-Utilities

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy

PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services
> Privacy Statement | Freedom of Information Act | Disclaimer

> > Oct 13 2004 06:44:09

h







Go

Entrez

PubMed

Nucleotide

otide Protein

Genoms

Structure

OMIM

PMO

Journals

Books

Search PubMed

Limits

for AL-2 AND vasculature

Preview/Index

History

Clipboard

Clear

Details

No items found.

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Book
Search	PubMed	fo	r EFL-6				Go	Clear	Book
		Limits	Previe	w/Index	History	Clipt	oard	Det	ails
About Entr	er	One of your to See <u>Details</u> .	erms is not	found in the	e database.				
Text Version	on	Display Sui	mmary		Show: 500	Sort		Send to	ext
Entrez Pu	bMed	:	Items 1 - 16	of 16	, , , , , , , , , , , , , , , , , , , ,				One p
Overview Help FAQ		□ 1: <u>Fav DS</u> ,	Qiu X, Large	E. Smith CP.	Mango S, Joha	nson BL.		Related A	rticles, Li
Tutorial New/Notew E-Utilities PubMed 8	·	35/Rb, Dev Bio	pha-1, and l. 2004 Jul 1;2	ubc-18. 271(1):11-25.	pharyngeal de		t in C. e	elegans by	lin-
Journals Date MeSH Date Single Citat		2: Dellaca Del	RL, Santus P. otti A, Calverl	Aliverti A. S ev PM	tevenson N. Cer	ntanni S, Mac	<u>klem</u>	Related Ar	ticles, Li
Batch Citat Clinical Qui LinkOut Cubby	ion Matcher	techniq Eur Resp	u e . oir J. 2004 Fel	0;23(2):232-4	imitation in (0. ed for MEDLIN		g the fo	orced oscill	lation
Related R		3: Baydur /	A. Wilkinson I	L. Mehdian R	. Bains B, Milic	o-Emili J.		Related Ar	ticles, [.j
Order Documents NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts		Extrathoracic expiratory flow limitation in obesity and obstructive and restrictive disorders: effects of increasing negative expiratory pressure. Chest. 2004 Jan;125(1):98-105. PMID: 14718427 [PubMed - indexed for MEDLINE]							
ClinicalTrial PubMed Ce		□ 4: Wang PI	I, Kuo PH, H	su CL, Wu Hl	D, Chang YS, K	tuo SH, Yang	rPC.	Related Ar	ticles, Li
		Diagno Chest, 20	stic value o 003 Nov;124(f negative 6 5):1762-7.	expiratory pre	essure for a			•
		5: Chen J. Laughlin LS, Hendrickx AG, Natarajan K. Overstreet JW, Lasley Related Articles, Li							
		The effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on chorionic gonadotrophin activity in pregnant macaques. Toxicology. 2003 Apr 15;186(1-2):21-31. PMID: 12604168 [PubMed - indexed for MEDLINE]							
		6: Boxem M	1, van den He	uvel S.				Related Art	icles, Li
		Curr Biol	l. 2002 Jun 4;]	12(11):906-11	ultivulva gen d for MEDLINE		(1) regu		
		7: Boni E. C	Corda L, Franc C	chini D, Chiro	<u>li P, Damiani G</u>	P, Pini L, Gr	<u>assi V.</u>	Related Art	icl e s, Li
		Volume	effect and	exertional o	lyspnoea afte	r bronchod	ilator ii	n patients	with

8: landelli I, Aliverti A, Kayser B, Dellaca R, Cala SJ, Duranti R, Kelly S, Scano G, Sliwinski P, Yan S, Macklem PT, Pedotti A.

b e

COPD with and without expiratory flow limitation at rest.

e ch

PMID: 12037229 [PubMed - indexed for MEDLINE]

Thorax. 2002 Jun;57(6):528-32.

fcg

cb

h g

e e

Determinants of exercise performance in normal men with externally imposed expiratory flow limitation. J Appl Physiol. 2002 May;92(5):1943-52. PMID: 11960944 [PubMed - indexed for MEDLINE] 9: Ferretti A. Giampiccolo P. Cavalli A. Milic-Emili J. Tantucci C. Related Articles, Li Expiratory flow limitation and orthopnea in massively obese subjects. Chest. 2001 May;119(5):1401-8. PMID: 11348945 [PubMed - indexed for MEDLINE] 10: Michaud S, Bourgault AM, Gaudreau C. Related Articles, Li Epidemiology of enterococcal bacteremia in a referral center for hepatobiliar diseases. Infection. 2000 Nov-Dec;28(6):361-6. PMID: 11139155 [PubMed - indexed for MEDLINE] 11: Kumar J, van Rheenen HA. Related Articles, Li A major gene for time of flowering in chickpea. J Hered. 2000 Jan-Feb; 91(1):67-8. PMID: 10739130 [PubMed - indexed for MEDLINE] 12: McClaran SR, Harms CA, Pegelow DF, Dempsey JA Related Articles, Li Smaller lungs in women affect exercise hyperpnea. J Appl Physiol. 1998 Jun;84(6):1872-81. PMID: 9609779 [PubMed - indexed for MEDLINE] 13: Peslin R, Farre R, Rotger M, Navajas D. Related Articles, Li Effect of expiratory flow limitation on respiratory mechanical impedance: a model study. J Appl Physiol. 1996 Dec;81(6):2399-406. PMID: 9018485 [PubMed - indexed for MEDLINE] 114: Barret J, Chitayat D, Sermer M, Amankwah K, Morrow R, Toi A, Ryan Related Articles, Li The prognostic factors in the prenatal diagnosis of the echogenic fetal lung. Prenat Diagn. 1995 Sep;15(9):849-53. PMID: 8559756 [PubMed - indexed for MEDLINE] 15: Yamazaki S. Related Articles, Li Fibrous structure of the joint capsule in the human shoulder. Okajimas Folia Anat Jpn. 1990 Aug;67(2-3):127-39. PMID: 2216306 [PubMed - indexed for MEDLINE] 16: Einarsdottir GH, Stankovich MT, Tu SC. Related Articles, Li Studies of electron-transfer properties of salicylate hydroxylase from Pseudomonas cepacia and effects of salicylate and benzoate binding. Biochemistry. 1988 May 3;27(9):3277-85. PMID: 3390431 [PubMed - indexed for MEDLINE] Display Summary Show: 500 Sort Send to

Write to the Help Desk

NCB! | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09

Related Articles, Li







of Medicine Entrez PubMed Nucleotide Protein Genome Structure MIMO PMC Journals Books Search PubMed * for ephrin-B3 Go Clear Limits Preview/Index History Clipboard Details Display Summary Show: |500 Send to Text About Entrez Items 1 - 41 of 41 One pa **Text Version** 1: Wieland I, Jakubiczka S, Muschke P, Cohen M, Thiele H, Gerlach KL. Related Articles, Li Adams RH, Wieacker P. Entrez PubMed Mutations of the ephrin-B1 gene cause craniofrontonasal syndrome. Overview Am J Hum Genet. 2004 Jun;74(6):1209-15. Epub 2004 Apr 29. Help | FAQ PMID: 15124102 [PubMed - indexed for MEDLINE] Tutorial New/Noteworthy 2: Bembenek ME, Schmidt S, Li P, Morawiak J, Prack A, Jain S, Roy R, Related Articles, Li E-Utilities Parsons T, Chee L. Characterization of the kinase domain of the ephrin-B3 receptor tyrosine kinase PubMed Services using a scintillation proximity assay. Journals Database MeSH Database Assay Drug Dev Technol. 2003 Aug;1(4):555-63. Single Citation Matcher PMID: 15090252 [PubMed - indexed for MEDLINE] **Batch Citation Matcher** Clinical Queries 3: Pascall JC, Brown KD. Related Articles, Li LinkOut Cubby Intramembrane cleavage of ephrinB3 by the human rhomboid family protease RHBDL2. Related Resources Biochem Biophys Res Commun. 2004 Apr 23;317(1):244-52. Order Documents PMID: 15047175 [PubMed - indexed for MEDLINE] **NLM Catalog NLM Gateway** 1 4: Chen ZY, Sun C, Reuhl K, Bergemann A, Henkemeyer M, Zhou R. Related Articles, Li TOXNET Abnormal hippocampal axon bundling in EphB receptor mutant mice. Consumer Health Clinical Alerts J Neurosci. 2004 Mar 10;24(10):2366-74. ClinicalTrials.gov PMID: 15014111 [PubMed - indexed for MEDLINE] PubMed Central 5: Grunwald IC, Korte M, Adelmann G, Plueck A, Kullander K, Adams Related Articles, Li RH, Frotscher M, Bonhoeffer T, Klein R Hippocampal plasticity requires postsynaptic ephrinBs. Nat Neurosci. 2004 Jan;7(1):33-40. Epub 2003 Dec 14. PMID: 14699416 [PubMed - indexed for MEDLINE] 6: Yu G, Luo H, Wu Y, Wu J. Related Articles, Li Mouse ephrinB3 augments T-cell signaling and responses to T-cell receptor ligation. J Biol Chem. 2003 Nov 21;278(47):47209-16. Epub 2003 Sep 17. PMID: 13679370 [PubMed - indexed for MEDLINE] 7: Egawa M, Yoshioka S, Higuchi T, Sato Y, Tatsumi K, Fujiwara H, Fujii Related Articles, Li Ephrin B1 is expressed on human luteinizing granulosa cells in corpora lutea (the early luteal phase: the possible involvement of the B class Eph-ephrin system during corpus luteum formation. J Clin Endocrinol Metab. 2003 Sep, 88(9): 4384-92. PMID: 12970314 [PubMed - indexed for MEDLINE]

developing mammalian spinal cord.
Prog Neurobiol. 2003 Jul;70(4):347-61. Review.
PMID: 12963092 [PubMed - indexed for MEDLINE]

Physiological, anatomical and genetic identification of CPG neurons in the

8: Kiehn O, Butt SJ.

		0
□9:	de Saint-Vis B, Bouchet C, Gautier G, Valladeau J, Caux C, Garrone P.	Related Articles, Li
	Human dendritic cells express neuronal Eph receptor tyrosine EphA2 in regulating adhesion to fibronectin. Blood. 2003 Dec 15;102(13):4431-40. Epub 2003 Aug 07. PMID: 12907451 [PubMed - indexed for MEDLINE]	e kinases: role of
□ 10	: Tabakoff B. Bhave SV, Hoffman PL.	Related Articles, Li
	Selective breeding, quantitative trait locus analysis, and gen candidate genes for complex drug-related behaviors. J Neurosci. 2003 Jun 1;23(11):4491-8. PMID: 12805289 [PubMed - indexed for MEDLINE]	e arrays identify
□11	Brors D, Bodmer D, Pak K, Aletsee C, Schafers M, Dazert S, Ryan AF.	Related Articles, Li
	EphA4 provides repulsive signals to developing cochlear gamediated through ephrin-B2 and -B3. J Comp Neurol. 2003 Jul 14;462(1):90-100. PMID: 12761826 [PubMed - indexed for MEDLINE]	nglion neurites
□ 12	Kubis N. Catala M.	Related Articles, Li
	[Development and maturation of the pyramidal tract] Neurochirurgie. 2003 May;49(2-3 Pt 2):145-53. Review. French. PMID: 12746689 [PubMed - indexed for MEDLINE]	
13 :	Sanchez-Carbayo M, Belbin TJ, Scotlandi K, Prystowsky M, Baldini N, Childs G, Cordon-Cardo C.	Related Articles, Li
	Expression profiling of osteosarcoma cells transfected with ligenes: regulation of cell adhesion, apoptosis, and tumor supplemes. Lab Invest. 2003 Apr;83(4):507-17. PMID: 12695554 [PubMed - indexed for MEDLINE]	MDR1 and NEO pression-related
□ 14:	Howard MA, Rodenas-Ruano A, Henkemeyer M, Martin GK, Lonsbury-Martin BL, Liebl DJ	Related Articles, Li
	Eph receptor deficiencies lead to altered cochlear function. Hear Res. 2003 Apr;178(1-2):118-30. PMID: 12684184 [PubMed - indexed for MEDLINE]	
□ 15:	Pickles JO.	Related Articles, Li
	Expression of Ephs and ephrins in developing mouse inner e Hear Res. 2003 Apr;178(1-2):44-51. PMID: 12684176 [PubMed - indexed for MEDLINE]	ar.
□16:	Kullander K, Butt SJ, Lebret JM, Lundfald L, Restrepo CE, Rydstrom A, Klein R, Kiehn O.	Related Articles, Li
	Role of EphA4 and EphrinB3 in local neuronal circuits that of Science. 2003 Mar 21;299(5614):1889-92. PMID: 12649481 [PubMed - indexed for MEDLINE]	control walking.
□ 17:	Bianchi LM, Dinsio K, Davoli K, Gale NW.	Related Articles, Li
	Lac z Histochemistry and immunohistochemistry reveal ephrexpression in the inner ear. J Histochem Cytochem. 2002 Dec;50(12):1641-5. PMID: 12486086 [PubMed - indexed for MEDLINE]	rin-B ligand
□ 18:	<u>Takemoto M, Fukuda T, Sonoda R, Murakami F, Tanaka H, Yamamoto N.</u>	Related Articles, Li
	Ephrin-B3-EphA4 interactions regulate the growth of specific axon populations in vitro. Eur J Neurosci. 2002 Sep;16(6):1168-72.	c thalamocortica

PMID: 12383247 [PubMed - indexed for MEDLINE]

19: Varelias A, Koblar SA, Cowled PA, Carter CD, Claver M.

Related Articles. Li

Human osteosarcoma expresses specific ephrin profiles: implications for tumorigenicity and prognosis.

Cancer. 2002 Aug 15;95(4):862-9.

PMID: 12209731 [PubMed - indexed for MEDLINE]

20: Hindges R, McLaughlin T, Genoud N, Henkemeyer M, O'Leary DD. Related Articles, Li

EphB forward signaling controls directional branch extension and arborization required for dorsal-ventral retinotopic mapping.

Neuron. 2002 Aug 1;35(3):475-87.

PMID: 12165470 [PubMed - indexed for MEDLINE]

21: Liu W, Ahmad SA, Jung YD, Reinmuth N, Fan F, Bucana CD, Ellis Related Articles, Li LM.

Coexpression of ephrin-Bs and their receptors in colon carcinoma. Cancer. 2002 Feb 15;94(4):934-9.

PMID: 11920461 [PubMed - indexed for MEDLINE]

22: Grunwald IC, Korte M, Wolfer D, Wilkinson GA, Unsicker K, Lipp Related Articles, Li HP, Bonhoeffer T, Klein R.

Kinase-independent requirement of EphB2 receptors in hippocampal synapti plasticity.

Neuron. 2001 Dec 20;32(6):1027-40.

PMID: 11754835 [PubMed - indexed for MEDLINE]

1 23: Coonan JR, Greferath U, Messenger J, Hartley L, Murphy M, Boyd Related Articles, Li AW, Dottori M, Galea MP, Bartlett PF.

Development and reorganization of corticospinal projections in EphA4 deficient mice.

J Comp Neurol. 2001 Jul 23;436(2):248-62.

PMID: 11438928 [PubMed - indexed for MEDLINE]

24: Stuckmann I, Weigmann A, Shevchenko A, Mann M, Huttner WB. Related Articles, Li

Ephrin B1 is expressed on neuroepithelial cells in correlation with neocortical neurogenesis.

J Neurosci. 2001 Apr 15;21(8):2726-37. PMID: 11306625 [PubMed - indexed for MEDLINE]

25: Kullander K, Croll SD, Zimmer M, Pan L, McClain J, Hughes V, Related Articles, Li Zabski S, DeChiara TM, Klein R, Yancopoulos GD, Gale NW

Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control.

Genes Dev. 2001 Apr 1;15(7):877-88.

PMID: 11297511 [PubMed - indexed for MEDLINE]

26: Rutherford A.

Related Articles, Li

Looking into mirror movement disorder.

Trends Mol Med. 2001 Mar;7(3):101. No abstract available.

PMID: 11286777 [PubMed - indexed for MEDLINE]

1727: Yokoyama N. Romero MI, Cowan CA, Galvan P, Helmbacher F. Related Articles, Li Charnay P, Parada LF, Henkemeyer M.

Forward signaling mediated by ephrin-B3 prevents contralateral corticospina axons from recrossing the spinal cord midline.

Neuron. 2001 Jan;29(1):85-97.

PMID: 11182083 [PubMed - indexed for MEDLINE]

Tang XX, Zhao H, Robinson ME, Cnaan A, London W, Cohn SL,

Related Articles, Li

h

Cheung NK, Brodeur GM, Evans AE, Ikegaki N.

Prognostic significance of EPHB6, EFNB2, and EFNB3 expressions in neuroblastoma.

Med Pediatr Oncol. 2000 Dec;35(6):656-8.

PMID: 11107140 [PubMed - indexed for MEDLINE]

1 29: Tang XX, Zhao H, Robinson ME, Cohen B, Chaan A, London W, Cohn Related Articles, Li SL, Cheung NK, Brodeur GM, Evans AE, Ikegaki N.



Implications of EPHB6, EFNB2, and EFNB3 expressions in human neuroblastoma.

Proc Natl Acad Sci U S A. 2000 Sep 26;97(20):10936-41. PMID: 10984508 [PubMed - indexed for MEDLINE]

30: L'Allemain G.

Related Articles, Li



[Involvement of ephrins and their receptors in oncogenesis]

Bull Cancer. 2000 Jul;87(7-8):529-30. French.

PMID: 10969208 [PubMed - indexed for MEDLINE]

31: Helbling PM, Saulnier DM, Robinson V, Christiansen JH, Wilkinson Related Articles, Li DG, Brandli AW.



Comparative analysis of embryonic gene expression defines potential interaction sites for Xenopus EphB4 receptors with ephrin-B ligands.

Dev Dyn. 1999 Dec;216(4-5):361-73.

PMID: 10633856 [PubMed - indexed for MEDLINE]

32: Helbling PM, Saulnier DM, Brandli AW.

Related Articles, Li



The receptor tyrosine kinase EphB4 and ephrin-B ligands restrict angiogenic growth of embryonic veins in Xenopus laevis.

Development. 2000 Jan; 127(2):269-78.

PMID: 10603345 [PubMed - indexed for MEDLINE]

33: van Heumen WR, Claxton C, Pickles JO.

Related Articles, Li



Expression of EphA4 in developing inner ears of the mouse and guinea pig. Hear Res. 2000 Jan;139(1-2):42-50.

PMID: 10601711 [PubMed - indexed for MEDLINE]

Tang XX, Evans AE, Zhao H, Chaan A, London W, Cohn SL, Brodeur Related Articles, Li GM, Ikegaki N.



High-level expression of EPHB6, EFNB2, and EFNB3 is associated with lov tumor stage and high TrkA expression in human neuroblastomas.

Clin Cancer Res. 1999 Jun;5(6):1491-6.

PMID: 10389937 [PubMed - indexed for MEDLINE]

T35: Miranda JD, White LA, Marcillo AE, Willson CA, Jagid J, Whittemore Related Articles, Li



Induction of Eph B3 after spinal cord injury.

Exp Neurol. 1999 Mar;156(1):218-22.

PMID: 10192794 [PubMed - indexed for MEDLINE]

☐ 36: O'Leary DD, Wilkinson DG.

Related Articles, Li



Eph receptors and ephrins in neural development.

Curr Opin Neurobiol. 1999 Feb;9(1):65-73. Review. PMID: 10072375 [PubMed - indexed for MEDLINE]

37: Tang XX, Brodeur GM, Campling BG, Ikegaki N.

Related Articles, Li



Coexpression of transcripts encoding EPHB receptor protein tyrosine kinases and their ephrin-B ligands in human small cell lung carcinoma.

Clin Cancer Res. 1999 Feb;5(2):455-60.

PMID: 10037197 [PubMed - indexed for MEDLINE]

38: Lin D, Gish GD, Songyang Z, Pawson T. Related Articles, Li The carboxyl terminus of B class ephrins constitutes a PDZ domain binding motif. J Biol Chem. 1999 Feb 5;274(6):3726-33. PMID: 9920925 [PubMed - indexed for MEDLINE] 39: Flanagan JG, Vanderhaeghen P. Related Articles, Li The ephrins and Eph receptors in neural development. Annu Rev Neurosci. 1998;21:309-45. Review. PMID: 9530499 [PubMed - indexed for MEDLINE] 1 40: Bergemann AD, Zhang L, Chiang MK, Brambilla R, Klein R, Flanagan Related Articles, Li Ephrin-B3, a ligand for the receptor EphB3, expressed at the midline of the developing neural tube. Oncogene. 1998 Jan 29;16(4):471-80. PMID: 9484836 [PubMed - indexed for MEDLINE] 41: Tang XX, Pleasure DE, Ikegaki N. Related Articles, Li cDNA cloning, chromosomal localization, and expression pattern of EPLG8, new member of the EPLG gene family encoding ligands of EPH-related protein-tyrosine kinase receptors. Genomics. 1997 Apr 1;41(1):17-24. PMID: 9126477 [PubMed - indexed for MEDLINE]

Write to the Help Desk
NCB! | NLM | NIH
Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer

◆ Show: 500 ★ Sort

Oct 13 2004 06:44:09

Send to


h

Display

Summary







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Book
		for					Go	Clear	
		Limits	Previe	10.1110050	History	Clip	poard	Deta	ils
About Entre	DZ	Display Abst	ract	*	Show: 20				xt

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher **Batch Citation Matcher** Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

☐ 1: Genomics. 1997 Apr 1;41(1):17-24.

Related Articles, I

TARRESTON SCHOOL FULL-TEXT ARTICLE

cDNA cloning, chromosomal localization, and expression pattern EPLG8, a new member of the EPLG gene family encoding ligand of EPH-related protein-tyrosine kinase receptors.

Tang XX, Pleasure DE, Ikegaki N.

Division of Neurology Research, Children's Hospital of Philadelphia, Pennsylvania 19104-4318, USA.

By screening a human fetal brain cDNA library under low stringency using cDNA encoding the mouse ligand of Cek5 as a probe, we have isolated a nove cDNA belonging to the EPLG gene family. This family encodes ligands of EP related tyrosine kinase receptors. Since the novel gene is the eighth member of the EPLG gene family, it is designated EPLG8. The deduced amino acid sequence of EPLG8 suggests that it encodes a transmembrane protein that is m related to those encoded by EPLG2 and EPLG5. We mapped the EPLG8 gene human chromosome 17p11.2-p13.1 by PCR screening of human-rodent somati cell hybrid panels. In the midterm fetus, EPLG8 mRNA is expressed at the highest level in brain, followed by heart, kidney, and lung. In the adult, EPLG mRNA expression is restricted to brain. These data suggest that LERK-8, the protein encoded by EPLG8, is important in brain development as well as in its maintenance. Moreover, since levels of EPLG8 expression were particularly him in several forebrain subregions compared to other brain subregions, LERK-8 n play a pivotal role in forebrain function.

PMID: 9126477 [PubMed - indexed for MEDLINE]

Display Abstract	Show:	20 💌	Sort •	Send to Text
------------------	-------	------	--------	--------------

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09

h

cb.

fcg c

e e

b e

b C







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	MIMO	PMC	Journals	Books
Search P	ubMed	for					Go	Clear	
		Limits	Previe	w/index	History	Clip	board	Deta	ils
About Entre:		Display Abs	tract		Show: 20	Sort		Send to Te	xt

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby

Related Resources Order Documents NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

□ 1: Oncogene. 1998 Jan 29;16(4):471-80.

species gnideleties surress and

Related Articles, I

Ephrin-B3, a ligand for the receptor EphB3, expressed at the midline of the developing neural tube.

Bergemann AD, Zhang L, Chiang MK, Brambilla R, Klein R, Flanagan J

Department of Cell Biology, Harvard Medical School, Boston, Massachusetts 02115, USA.

The ephrins are a family of ligands that bind to Eph family receptor tyrosine kinases, and have been implicated in axon guidance and other patterning processes during vertebrate development. We describe here the identification ϵ characterization of murine ephrin-B3. The cDNA encodes a 340 amino acid transmembrane molecule, most closely related to the two other known transmembrane ligands, ephrin-B1 and ephrin-B2. In addition to homology in their extracellular receptor binding domains, these transmembrane ligands shar striking homology between their cytoplasmic domains, with 31 of the last 34 amino acids of ephrin-B3 being identical to ephrin-B2, suggesting functional interactions of the cytoplasmic tail. While most Eph family ligands are promiscuous in their interactions with Eph receptors, binding studies with the: receptors known to bind other transmembrane ligands only revealed a high affinity interaction of ephrin-B3 with EphB3, with a dissociation constant of approximately 1 nM. In situ hybridization of mouse embryos showed ephrin-B is expressed prominently at the dorsal and ventral midline of the neural tube, particularly in the floor plate, a structure with key functions in patterning the nervous system. The isolation of this ligand may help to elucidate the molecula basis of patterning activities at the neural tube midline.

PMID: 9484836 [PubMed - indexed for MEDLINE]

Display Abstract		00	10-4	
Abstract *	Show:	20 *	ISOπ SEE	Send to Text
		Section of the sectio	Secretaria de la constante de	

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13/2004/06:44:09







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Book
Search P	ubMed	for					Go	Clear	
		Limits	Previe	w/Index	History	Clip	board	Deta	ails
About Caters						Sort		• • • • • • • •	ext

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
Clinical Trials.gov
PubMed Central

h

1: Exp Neurol. 1999 Mar; 156(1):218-22.

Related Articles, L

FULL-TEXT ARTICLE

Induction of Eph B3 after spinal cord injury.

Miranda JD, White LA, Marcillo AE, Willson CA, Jagid J, Whittemore S.

Department of Neurological Surgery, University of Miami School of Medicine 1600 Northwest 10th Avenue, R-48, Miami, Florida 33136, USA.

Spinal cord injury (SCI) in adult rats initiates a cascade of events producing a nonpermissive environment for axonal regeneration. This nonfavorable environment could be due to the expression of repulsive factors. The Eph recel protein tyrosine kinases and their respective ligands (ephrins) are families of molecules that play a major role in axonal pathfinding and target recognition during central nervous system (CNS) development. Their mechanism of action mediated by repellent forces between receptor and ligand. The possible role the these molecules play after CNS trauma is unknown. We hypothesized that an increase in the expression of Eph proteins and/or ephrins may be one of the molecular cues that restrict axonal regeneration after SCI. Rats received a contusive SCI at T10 and in situ hybridization studies 7 days posttrauma demonstrated: (i) a marked up-regulation of Eph B3 mRNA in cells located in white matter at the lesion epicenter, but not rostral or caudal to the injury site, (ii) an increase in Eph B3 mRNA in neurons in the ventral horn and intermedia zone of the gray matter, rostral and caudal to the lesion. Immunohistochemical analyses localizing Eph B3 protein were consistent with the mRNA results. Colocalization studies performed in injured animals demonstrated increased E_l B3 expression in white matter astrocytes and motor neurons of the gray matter These results suggest that Eph B3 may contribute to the unfavorable environm for axonal regeneration after SCI. Copyright 1999 Academic Press.

PMID: 10192794 [PubMed - indexed for MEDLINE]

District Abetreet					
Show: 20 * Sort * Send to Text	Display Abstract	a 20	Sort	al Consti	

Write to the Help Desk

NCB! | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09

C







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Books
Search	PubMed	for		***************************************			Go	Clear	
		Limits	Previe	w/Index	History	Clip	board	Details	;
ihord Cat	(A)	Display Abst	ract		a. 20 8	Sort		Send to Text	

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Malcher
Clinical Queries
LinkOut
Cubby

Related Resources Order Documents NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central **1:** Genes Dev. 2001 Apr 1;15(7):877-88.

Related Articles, I

FREE full text article of www.genesdev.org

Ephrin-B3 is the midline barrier that prevents corticospinal tract axons from recrossing, allowing for unilateral motor control.

Kullander K, Croll SD, Zimmer M, Pan L, McClain J, Hughes V, Zabski DeChiara TM, Klein R, Yancopoulos GD, Gale NW.

European Molecular Biology Laboratory, D-69117 Heidelberg, Germany.

Growing axons follow highly stereotypical pathways, guided by a variety of attractive and repulsive cues, before establishing specific connections with dist targets. A particularly well-known example that illustrates the complexity of axonal migration pathways involves the axonal projections of motor neurons located in the motor cortex. These projections take a complex route during whi they first cross the midline, then form the corticospinal tract, and ultimately connect with motor neurons in the contralateral side of the spinal cord. These obligatory contralateral connections account for why one side of the brain controls movement on the opposing side of the body. The netrins and slits prov well-known midline signals that regulate axonal crossings at the midline. Here we report that a member of the ephrin family, ephrin-B3, also plays a key role the midline to regulate axonal crossing. In particular, we show that ephrin-B3; as the midline barrier that prevents corticospinal tract projections from recross: when they enter the spinal gray matter. We report that in ephrin-B3(-/-) mice, corticospinal tract projections freely recross in the spinal gray matter, such tha the motor cortex on one side of the brain now provides bilateral input to the sp cord. This neuroanatomical abnormality in ephrin-B3(-/-) mice correlates with loss of unilateral motor control, yielding mice that simultaneously move their right and left limbs and thus have a peculiar hopping gait quite unlike the alternate step gait displayed by normal mice. The corticospinal and walking defects in ephrin-B3(-/-) mice resemble those recently reported for mice lackir. the EphA4 receptor, which binds ephrin-B3 as well as other ephrins, suggestin that the binding of EphA4-bearing axonal processes to ephrin-B3 at the midlin provides the repulsive signal that prevents corticospinal tract projections from recrossing the midline in the developing spinal cord.

PMID: 11297511 [PubMed - indexed for MEDLINE]

Display Abstract She	ow: 20 • Sort • Send to Text
----------------------	------------------------------

Write to the Help Desk

C







Entrez	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Books
Search P	ubMed	for					Go	Clear	
		Limits	Previe	w/Index	History	Clip	board	Detail	S
Ahout Sotra:	,	Display Ahe	tract		- 120	Sort	*	Send to Tex	tt

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.goy PubMed Central

1: Eur J Neurosci. 2002 Sep;16(6):1168-72.

Related Articles, I



Ephrin-B3-EphA4 interactions regulate the growth of specific thalamocortical axon populations in vitro.

Takemoto M, Fukuda T, Sonoda R, Murakami F, Tanaka H, Yamamoto I

Neuroscience Laboratories, Graduate School of Frontier Biosciences, Osaka University, Toyonaka, Osaka 560-8531, Japan.

The role was studied of ephrin-B3, a ligand of the Eph family of tyrosine kinas receptors, in the formation of cortical connectivity. In situ hybridization and immunohistochemistry showed that EphA4, a receptor of ephrin-B3, was expressed in the lateral thalamus (visual and somaotosensory thalamus) of the developing rat brain, but not in the medial thalamic nuclei which project to the limbic cortex. Correspondingly, ephrin-B3 was expressed strongly in the developing limbic cortex including amygdala, entorhinal cortex and hippocampus. To examine the action of ephrin-B3 on thalamic axons, either lateral or medial thalamic explants were cultured on membranes obtained from ephrin-B3-expressing COS cells. Axonal growth was inhibited for cells from the lateral thalamus but not from the medial thalamus. These results suggest that ephrin-B3 contributes to regional specificity by suppressing axonal growth of lateral thalamic neurons.

PMID: 12383247 [PubMed - indexed for MEDLINE]

Display Abstract - Sh	how: 20 💌	Sort -	Send to Text
-----------------------	-----------	--------	--------------

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09

h

cb


h g e fcg c

b

b C







				79					
Entrez	PubMed	Nucleotide	Protein	Genome	Structure	MIMO	PMC	Journals	Books
Search	PubMed	for				***************************************	Go	Clear	
		Limits	Previe	w/Index	History	Clip	phoard	Detai	ls
About Entr	ez	Display Abs	tract		Show: 20	Sort		end to Te	đ
Text Versio	ori	□ 1: J Biol Cl	nem. 2003	Nov 21;27	8(47):47209-	16. Epub	2003 Sep	Relat	ed Arti
		1/.							į

Entrez PubMed Overview Help | FAQ Tutorial. New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher **Batch Citation Matcher** Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Mouse ephrinB3 augments T-cell signaling and responses to T-cel receptor ligation.

Yu G, Luo H, Wu Y, Wu J.

FRIE full text article at

www.jbc.org

Laboratory of Immunology and the Nephrology Service of Notre Dame Hospit Centre Hospitalier de l'Universite de Montreal, Universite de Montreal, Montre Quebec H2L 4M1, Canada.

Ephrins (EFN) are cell-surface ligands of Ephs, the largest family of cell-surfa receptor tyrosine kinases. The function of EFNs in the immune system has not been well studied, although some EFNs and Ephs are expressed at high levels certain leukocytes. We report here that EFNB3 and its receptors (collectively called EFNB3Rs, as EFNB3 binds to multiple EphBs) were expressed in peripheral T cells and monocytes/macrophages, with T cells being the dominal EFNB3+ and EFNB3R+ cell type. Solid-phase EFNB3-Fc in the presence of suboptimal anti-CD3 crosslinking enhanced T-cell responses in terms of proliferation, activation marker expression, interferon-gamma but not interleuk 2 production, and cytotoxic T-cell activity. EFNB3R costimulation in the presence of phorbol 12-myristate 13- acetate was insensitive to cyclosporin A, similar to CD28 costimulation, suggesting they might share a part of the signal pathway. After crosslinking, T-cell receptor and EFNB3R congregated into aggregated rafts, and this provided a morphological basis for signaling pathwa of T-cell receptor and EFNB3R to interact. Solid-phase EFNB3-Fc augmented p38 and p44/42 MAPK activation further downstream of the signaling pathwar These data suggest that EFNB3 is important in T-cell/T-cell and T-cell/antiger. presenting cell collaboration to enhance T-cell activation and function.

PMID: 13679370 [PubMed - indexed for MEDLINE]

fcg c

Display Abstract	Show: 20 •	Sort	Send to Text

Write to the Help Desk NOBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

b

b e

b

C

e e

Oct 13/2004/06:44:09

h g







Entrez	PubMed	Nucleotide	Protein	Genoms	Structure	MIMO	PMC	Journals	Books
Search P	ubMed	for				_	Go	Clear	
		Limits	Previe	w/Index	History	Clip	board	Deta	ails
Ahoui Entre:	7	Display Abst	ract		Show 20	Sort		A	ext

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher **Batch Citation Matcher** Clinical Queries LinkOut Cubby

Related Resources Order Documents **NLM Catalog** NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

1: J Comp Neurol. 2003 Jul 14;462(1):90-100.

Related Articles, I



EphA4 provides repulsive signals to developing cochlear ganglion neurites mediated through ephrin-B2 and -B3.

Brors D, Bodmer D, Pak K, Aletsee C, Schafers M, Dazert S, Ryan AF.

Department of Surgery, Division Otolaryngology and Neurosciences, Universi of California, San Diego School of Medicine and Veterans Administration Medical Center, La Jolla 92093, USA.

The ephrins and Eph receptors make up two large families of bi-directional signaling molecules that are known to play a role in the development of the nervous system. Recently, expression of EphA4 in the developing cochlea was shown, with strong expression in cells lining the osseous spiral lamina (OSL) through which afferent dendrites must pass to reach the organ of Corti (OC). It was also demonstrated that ephrin-B2 and -B3, both of which are known to interact with EphA4, are expressed by spiral ganglion (SG) neurons. To investigate the functional role of EphA4 in the development of inner ear neuro neonatal rat SG explants were cultured for 72 hours on uniformly coated surface or near stripes of EphA4/IgG-Fc-chimera. Control explants were cultured on o near IgG-Fc and EphA1/IgG-Fc-chimera. To assess the roles of ephrin-B2 and B3 in EphA4 signaling, SG explants were cultured with or without anti-ephrin and/or -B3 blocking antibodies. Growth patterns of SG neurites at the border o EphA4 receptor stripes showed repulsion, characterized by turning, stopping and/or reversal. In the case of IgG-Fc and EphA1, the neurites grew straight or the stripes. Treatment with either anti-ephrin-B2 or -B3 blocking antibodies significantly reduced the repulsive effect of an EphA4 stripe. Moreover, when both antibodies were used together, neurites crossed onto EphA4 stripes with r evidence of repulsion. The results suggest that EphA4 provides repulsive signa to SG neurites in the developing cochlea, and that ephrin-B2 and -B3 together mediate this response. Copyright 2003 Wiley-Liss, Inc.

PMID: 12761826 [PubMed - indexed for MEDLINE]

Display Abstract	Show: 20 Sort	Send to Text
Abstract -	Show: 20 Solt	- CAL

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

C

cb h g e e fcg c e e e b b e b

Related Articles, Li







of Medicine NIN Entrez PubMed Nucleotide Protein Genoms MIMO Structure PMC Journals Books Search | PubMed for lephrin AND vascularization Clear Go Limits Preview/Index History Clipboard Details Display Summary Show: |500 * Text About Entrez Items 1 - 35 of 35 One pa Text Version 1: Umeda N. Ozaki H. Hayashi H. Oshima K. Related Articles, Li Expression of ephrinB2 and its receptors on fibroproliferative membranes in Entrez PubMed ocular angiogenic diseases. Overview Am J Ophthalmol. 2004 Aug;138(2):270-9. Help | FAQ PMID: 15289137 [PubMed - indexed for MEDLINE] Tutorial New/Noteworthy E-Utilities 2: Liu W, Jung YD, Ahmad SA, McCarty MF, Stoeltzing O, Reinmuth N, Related Articles, Li Fan F, Ellis LM PubMed Services Effects of overexpression of ephrin-B2 on tumour growth in human colorectal Journals Database MeSH Database Br J Cancer. 2004 Apr 19;90(8):1620-6. Single Citation Matcher PMID: 15083195 [PubMed - indexed for MEDLINE] Batch Citation Matcher Clinical Queries 3: Noren NK, Lu M, Freeman AL, Koolpe M, Pasquale EB LinkOut Related Articles, Li Cubby Interplay between EphB4 on tumor cells and vascular ephrin-B2 regulates turn growth. Related Resources Proc Natl Acad Sci U S A. 2004 Apr 13;101(15):5583-8. Epub 2004 Apr 05. Order Documents PMID: 15067119 [PubMed - indexed for MEDLINE] **NLM Catalog NLM Gateway** 4: Patan S. TOXNET Related Articles, Li Consumer Health Vasculogenesis and angiogenesis. Clinical Alerts Cancer Treat Res. 2004;117:3-32. Review. ClinicalTrials.gov PubMed Central PMID: 15015550 [PubMed - indexed for MEDLINE] 5: Huang J, Soffer SZ, Kim ES, McCrudden KW, Huang J, New T, Manley Related Articles, Li CA, Middlesworth W, O'Toole K, Yamashiro DJ, Kandel JJ. Vascular remodeling marks tumors that recur during chronic suppression of angiogenesis. Mol Cancer Res. 2004 Jan;2(1):36-42. PMID: 14757844 [PubMed - indexed for MEDLINE] 6: Bicknell R, Harris AL Related Articles, Li Novel angiogenic signaling pathways and vascular targets. Annu Rev Pharmacol Toxicol. 2004;44:219-38. Review. PMID: 14744245 [PubMed - indexed for MEDLINE] 7: Sawai Y, Tamura S, Fukui K, Ito N, Imanaka K, Saeki A, Sakuda S, Kiso Related Articles, Li S, Matsuzawa Y. Expression of ephrin-B1 in hepatocellular carcinoma: possible involvement in neovascularization. J Hepatol. 2003 Dec;39(6):991-6. PMID: 14642617 [PubMed - indexed for MEDLINE]

VEGF-receptor inhibitors for anti-angiogenesis.
Nippon Yakurigaku Zasshi. 2003 Dec;122(6):498-503. Review.
PMID: 14639004 [PubMed - indexed for MEDLINE]

e ch

bе

8: Shibuya M.

h

cb

hg e e

e fcg

e ch

b e

		= -8- = 01 .
□9:	Augustin HG, Reiss Y.	Related Articles, Li
	EphB receptors and ephrinB ligands: regulators of vascular a homeostasis.	ssembly and
	Cell Tissue Res. 2003 Oct;314(1):25-31. Epub 2003 Aug 02. Review. PMID: 12905065 [PubMed - indexed for MEDLINE]	
1 0	: Sullivan DC, Bicknell R.	Related Articles, Li
	New molecular pathways in angiogenesis. Br J Cancer. 2003 Jul 21;89(2):228-31. Review. PMID: 12865906 [PubMed - indexed for MEDLINE]	
□ 11	: Andres AC, Munarini N, Djonov V, Bruneau S, Zuercher G, Loercher S, Rohrbach V, Ziemiecki A.	Related Articles, Li
	EphB4 receptor tyrosine kinase transgenic mice develop gloreminiscent of aglomerular vascular shunts. Mech Dev. 2003 Apr;120(4):511-6. PMID: 12676328 [PubMed - indexed for MEDLINE]	omerulopathies
□ 12	: Marme D.	Related Articles, Li
	VEGFs, angiopoietins, Ephrins and their receptors: putative	targets for tumo
₩	therapy? Ann Hematol. 2002;81 Suppl 2:S66. No abstract available. PMID: 12611080 [PubMed - indexed for MEDLINE]	
□ 13	: Augustin HG.	Related Articles, Li
	[Angiogenesis researchquo vadis?] Ophthalmologe. 2003 Feb;100(2):104-10. Review. German. PMID: 12589453 [PubMed - indexed for MEDLINE]	
1 4	Takahashi T, Takahashi K, St John PL, Fleming PA, Tomemori T, Watanabe T, Abrahamson DR, Drake CJ, Shirasawa T, Daniel TO.	Related Articles, Li
	A mutant receptor tyrosine phosphatase, CD148, causes defe	ects in vascular
-	development. Mol Cell Biol. 2003 Mar;23(5):1817-31. PMID: 12588999 [PubMed - indexed for MEDLINE]	
□ 15:	Straume O, Jackson DG, Akslen LA.	Řelated Articles, Li
	Independent prognostic impact of lymphatic vessel density a low-grade lymphangiogenesis in cutaneous melanoma. Clin Cancer Res. 2003 Jan;9(1):250-6. PMID: 12538477 [PubMed - indexed for MEDLINE]	and presence of
□16:	Kaban K, Herbst RS.	Related Articles, Li
	Angiogenesis as a target for cancer therapy. Hematol Oncol Clin North Am. 2002 Oct;16(5):1125-71. Review. PMID: 12512387 [PubMed - indexed for MEDLINE]	
□ 17:	Meyer S. Hafner C. Vogt T.	Related Articles, Li
65532 33533 33533	[Role of receptor tyrosine kinase in the angiogenesis] Hautarzt. 2002 Sep;53(9):629-42. Review. German. No abstract available PMID: 12432901 [PubMed - indexed for MEDLINE]	2 .
□ 18:	Brantley DM, Cheng N, Thompson EJ, Lin Q, Brekken RA, Thorpe PE, Muraoka RS, Cerretti DP, Pozzi A, Jackson D, Lin C, Chen J.	Related Articles, Li
	Soluble Eph A receptors inhibit tumor angiogenesis and prog Oncogene. 2002 Oct 10;21(46):7011-26. PMID: 12370823 [PubMed - indexed for MEDLINE]	ression in vivo.
19:	Papoutsi M, Othman-Hassan K, Christ B, Patel K, Wilting J.	Related Articles, Li

h

cb

h g

e fcg

e ch

b e

		1 450 3 01 4
	Development of an arterial tree in C6 gliomas but not in A2 Histochem Cell Biol. 2002 Sep;118(3):241-9. Epub 2002 Jul 13. PMID: 12271360 [PubMed - indexed for MEDLINE]	375 melanomas.
□ 20	: Nakamoto M, Bergemann AD.	Related Articles, Li
	Diverse roles for the Eph family of receptor tyrosine kinase Microsc Res Tech. 2002 Oct 1;59(1):58-67. Review. PMID: 12242697 [PubMed - indexed for MEDLINE]	es in carcinogenes
□ 21	: Steinle JJ, Meininger CJ, Forough R, Wu G, Wu MH, Granger HJ	Related Articles, Li
	Eph B4 receptor signaling mediates endothelial cell migrativia the phosphatidylinositol 3-kinase pathway. J Biol Chem. 2002 Nov 15;277(46):43830-5. Epub 2002 Sep 13. PMID: 12235151 [PubMed - indexed for MEDLINE]	on and proliferati
□ 22	: Straume O. Akslen LA.	Related Articles, Li
	Importance of vascular phenotype by basic fibroblast growth influence of the angiogenic factors basic fibroblast growth growth factor receptor-1 and ephrin-A1/EphA2 on melanon Am J Pathol. 2002 Mar;160(3):1009-19. PMID: 11891198 [PubMed - indexed for MEDLINE]	factor/fibroblast
□ 23	Cheng N. Brantley DM, Chen J.	Related Articles, Li
	The ephrins and Eph receptors in angiogenesis. Cytokine Growth Factor Rev. 2002 Feb;13(1):75-85. Review. PMID: 11750881 [PubMed - indexed for MEDLINE]	
□ 24	Kuwano M, Fukushi J, Okamoto M, Nishie A, Goto H, Ishibashi T, Ono M.	Related Articles, Li
	Angiogenesis factors. Intern Med. 2001 Jul;40(7):565-72. Review. PMID: 11506294 [PubMed - indexed for MEDLINE]	
□ 25	Gale NW, Baluk P, Pan L, Kwan M, Holash J, DeChiara TM, McDonald DM, Yancopoulos GD.	Related Articles, Li
	Ephrin-B2 selectively marks arterial vessels and neovascula adult, with expression in both endothelial and smooth-muscle Dev Biol. 2001 Feb 15;230(2):151-60. PMID: 11161569 [PubMed - indexed for MEDLINE]	rization sites in the cells.
□ 26:	Shin D. Garcia-Cardena G. Hayashi S. Gerety S. Asahara T. Stavrakis G. Isner J. Folkman J. Gimbrone MA Jr. Anderson DJ	Related Articles, Li
	Expression of ephrinB2 identifies a stable genetic difference and venous vascular smooth muscle as well as endothelial consubsets of microvessels at sites of adult neovascularization. Dev Biol. 2001 Feb 15;230(2):139-50. PMID: 11161568 [PubMed - indexed for MEDLINE]	e between arterial ells, and marks
□ 27:	Ogawa K. Pasqualini R. Lindberg RA, Kain R. Freeman AL, Pasquale EB	Related Articles, Li
	The ephrin-A1 ligand and its receptor, EphA2, are expressed neovascularization. Oncogene. 2000 Dec 7;19(52):6043-52. PMID: 11146556 [PubMed - indexed for MEDLINE]	l during tumor
□ 28:	Umess LD, Sorensen LK, Li DY.	Related Articles, Li
	Arteriovenous malformations in mice lacking activin receptor Nat Genet. 2000 Nov;26(3):328-31. PMID: 11062473 [PubMed - indexed for MEDLINE]	or-like kinase-1.

□ 29:	Yancopoulos GD, Davis S, Gale NW, Rudge JS, Wiegand SJ, Holash J.	Related Articles, Li
	Vascular-specific growth factors and blood vessel formation Nature. 2000 Sep 14;407(6801):242-8. Review. PMID: 11001067 [PubMed - indexed for MEDLINE]	
□30:	Yuan K, Jin YT, Lin MT.	Related Articles, Li
	Expression of Tie-2, angiopoietin-1, angiopoietin-2, ephrinE pyogenic granuloma of human gingiva implicates their roles angiogenesis. J Periodontal Res. 2000 Jun;35(3):165-71. PMID: 10929871 [PubMed - indexed for MEDLINE]	32 and EphB4 in in inflammatory
□31:	Myers C, Charboneau A, Boudreau N.	Related Articles, Li
	Homeobox B3 promotes capillary morphogenesis and angiog J Cell Biol. 2000 Jan 24;148(2):343-51. PMID: 10648567 [PubMed - indexed for MEDLINE]	genesis.
□ 32:	Easty DJ, Hill SP, Hsu MY, Fallowfield ME, Florenes VA, Herlyn M, Bennett DC	Related Articles, Li
	Up-regulation of ephrin-A1 during melanoma progression. Int J Cancer. 1999 Oct 22;84(5):494-501. PMID: 10502726 [PubMed - indexed for MEDLINE]	
□33:	Adams RH, Wilkinson GA, Weiss C, Diella F, Gale NW, Deutsch U, Risau W, Klein R.	Related Articles, Li
	Roles of ephrinB ligands and EphB receptors in cardiovascul demarcation of arterial/venous domains, vascular morphogen sprouting angiogenesis. Genes Dev. 1999 Feb 1;13(3):295-306. PMID: 9990854 [PubMed - indexed for MEDLINE]	ar development nesis, and
□34:	Chiarugi V, Magnelli L, Gallo O.	Related Articles, Li
	Cox-2, iNOS and p53 as play-makers of tumor angiogenesis Int J Mol Med. 1998 Dec;2(6):715-9. Review. PMID: 9850741 [PubMed - indexed for MEDLINE]	(review)
□ 35:	Pandey A, Shao H, Marks RM, Polverini PJ, Dixit VM.	Related Articles, Li
4:::::	Role of B61, the ligand for the Eck receptor tyrosine kinase, induced angiogenesis. Science. 1995 Apr 28;268(5210):567-9. PMID: 7536959 [PubMed - indexed for MEDLINE]	in TNF-alpha-
Display	Summary ★ Show: 500 ★ Sort ★ Se	nd to Text

Write to the Help Desk

NCB! | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Oct 13 2004 06:44:09